

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 1/04, 11/02, 29/16, 31/125, 303/24, 305/06, 305/10, 51/16, C11D 1/14		A1	(11) International Publication Number: WO 97/38956
			(43) International Publication Date: 23 October 1997 (23.10.97)
(21) International Application Number: PCT/US97/06339		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 16 April 1997 (16.04.97)		Published <i>With international search report.</i>	
(30) Priority Data: 60/015,523 16 April 1996 (16.04.96) US			
(71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).			
(72) Inventor; and (73) Inventor/Applicant (for US only): CONNOR, Daniel, Stedman [US/US]; 9217 Sagemeadow Drive, Cincinnati, OH 45251 (US).			
(74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).			
(54) Title: PROCESS FOR THE MANUFACTURE OF SURFACTANTS CONTAINING BRANCH CHAIN ALKYL GROUPS			
(57) Abstract Branched-chain olefins are prepared and used as feedstocks in the manufacture of deterative surfactants.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BV	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NR	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroun	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WO 97/38956

PCT/US97/06339

PROCESS FOR THE MANUFACTURE OF SURFACTANTS CONTAINING BRANCH CHAIN ALKYL GROUPS

FIELD OF THE INVENTION

5 The present invention relates to processes for manufacturing deterative surfactants, especially those containing branched-chain hydrophobic units.

BACKGROUND OF THE INVENTION

10 Conventional deterative surfactants comprise molecules having a water-solubilizing substituent (hydrophilic group) and an oleophilic substituent (hydrophobic group). Such surfactants typically comprise hydrophilic groups such as carboxylate, sulfate, sulfonate, amine oxide, polyoxyethylene, and the like, attached to an alkyl, alkenyl or alkaryl hydrophobe usually containing from about 10 to about 20 carbon atoms. Accordingly, the manufacturer of such surfactants must have access to a source of hydrophobe groups to which the desired hydrophile can be attached by chemical means. The earliest source of hydrophobe groups comprised the natural fats and oils, which were converted into soaps (i.e., carboxylate hydrophile) by saponification with base. Coconut oil and palm oil are still used to manufacture soap, as well as to manufacture the alkyl sulfate ("AS") class of surfactants. Other hydrophobes are available from petrochemicals, including 20 alkylated benzene which is used to manufacture alkyl benzene sulfonate surfactants ("LAS").

25 The literature asserts that certain branched hydrophobes can be used to advantage in the manufacture of alkyl sulfate deterative surfactants; see, for example, U.S. 3,480,556 to deWitt, et al., November 25, 1969. However, it has been determined that the beta-branched surfactants described in the '556 patent are inferior with respect to certain solubility parameters, as evidenced by their Krafft temperatures. It has further been determined that surfactants having branching towards the center of carbon chain of the hydrophobe have much lower Krafft temperatures. See: "The Aqueous Phase Behavior of Surfactants", R.G. Laughlin, 30 Academic Press, N.Y. (1994) p. 347. Accordingly, it has now been determined that such surfactants are preferred for use especially under cool or cold water washing conditions (e.g., 20°C-5°C).

35 One problem associated with the manufacture of deterative surfactants having hydrophobe groups with mid- or near-mid chain branching is the lack of a ready source of such hydrophobes. By the present invention, a process is described for manufacturing such branched hydrophobes and converting them into mid- or near-mid chain branched surfactants.

WO 97/38956

2

PCT/US97/06339

SUMMARY OF THE INVENTION

The present invention encompasses a process for preparing mid- to near mid-chain branched olefins (primarily, methyl branched at or near the mid-chain region). Such materials are then used as the basic feedstock which provides the hydrophobic portion of branched-chain deterative surfactants.

The process herein is designed to provide branched reaction products which are primarily (85%, or greater) alpha-olefins, and which are then converted into hydrophobes in the Oxo-reaction sequence noted hereinafter. Preferably, such branched alpha-olefins contain from about 11 to about 18 (avg.) total carbon atoms and comprise a linear chain having an average length in the 10-18 region. The branching is predominantly mono-methyl, but some di-methyl and some ethyl branching may occur. Advantageously, the present process results in little (1%, or less) geminal branching, i.e., little, if any, "quaternary" carbon substitution. Moreover, little (less than about 20%) vicinal branching occurs. Of course, some (ca. 20%) of the overall feedstock used in the subsequent Oxo-process may remain unbranched. Typically, and preferably from the standpoint of cleaning performance and biodegradability, the present process provides alpha-olefins with: an average number of branches (longest chain basis) in the 0.4-2.5 range; of the branched material, there are essentially no branches on carbons 1, 2 or on the terminal (omega) carbon of the longest chain of the branched material.

Following the formation and purification of the branched-chain alpha-olefin, the feedstock is subjected to an Oxo carbonylation process. In this Oxo-step, a catalyst (e.g., conventional cobalt carbonyl; see Kirk Othmer, below) which does not move the double bond from its initial position is used. This avoids the formation of vinylidene intermediates (which ultimately yield less favorable surfactants) and allows the carbonylation to proceed at the #1 and #2 carbon atoms.

All percentages, ratios and proportions herein are by weight, unless otherwise specified. All documents cited herein are, in relevant part, incorporated herein by reference.


DETAILED DESCRIPTION OF THE INVENTION

As can be seen from the foregoing, the present invention thus encompasses, in a process for preparing surfactant precursor hydrophobes from hydrocarbon feedstocks by the conversion of a coal or other hydrocarbon source to a mixture of carbon monoxide and hydrogen and the subsequent conversion of the carbon monoxide and hydrogen into a mixture of linear and branched hydrocarbons, the improvement which comprises abstracting from said mixture of linear and branched hydrocarbons the sub-set of branched hydrocarbons of the general formula:

WO 97/38956


PCT/US97/06339

3



wherein x is at least about 2, y is greater than or equal to 0 and wherein the sum of x+y is at least about 7.


5 The invention also encompasses, in a process for preparing surfactant precursor hydrophobes from coal or other hydrocarbon feedstocks by the conversion of such feedstocks to a mixture of carbon monoxide and hydrogen and the subsequent conversion of the carbon monoxide and hydrogen into a mixture of linear and branched hydrocarbons, the improvement which comprises abstracting from said mixture of linear and branched hydrocarbons the sub-set of branched hydrocarbons of
10 the general formula:




wherein p is at least about 2, q is 1 to 12, r is greater than or equal to 0 and the sum of p, q and r is at least about 6.

15 The invention also encompasses, in a process for preparing surfactant precursor hydrophobes from coal or other hydrocarbon feedstocks by the conversion of such feedstocks to a mixture of carbon monoxide and hydrogen and the subsequent conversion of the carbon monoxide and hydrogen into a mixture of linear and branched hydrocarbons, the improvement which comprises abstracting from said mixture of linear and branched hydrocarbons the set of branched hydrocarbons
20 comprising a mixture of:

(a) the sub-set of mono-methyl branched compounds of the formula:



(b) the sub-set of di-methyl branched compounds of the formula:



25 The foregoing branched hydrophobes can then be converted into the corresponding branched-chain deterative surfactants, in the manner disclosed hereinafter.

Process

30 Synthesis gas (carbon monoxide/hydrogen) can be produced from coal or other hydrocarbon feedstocks such as natural gas and used to build-up various

WO 97/38956

4

PCT/US97/06339

saturated and unsaturated linear, branched and cyclic hydrocarbons using conventional Fischer-Tropsch (F-T) chemistry. Such processes can be used to make a range of hydrocarbons to meet the gasoline, diesel and jet fuel needs. Two points with regard to the present invention are: first, recognition that branching occurs in F-T chemistry through free radical, not carbonium ion chemistry. This leads to isolated methyl branches with no gem-dimethyl, little ethyl and low levels of vicinal-dimethyl branches. Low pressure/low temp (i.e. wax producing) F-T chemistry builds up methylenes mostly in a linear fashion with typically about 1 methyl branch per 50 carbons. At higher pressures and/or higher temperatures (such as used for gasoline production) 1 methyl branch per 8 carbon atoms can be achieved. The rearrangement to form the methyl branch, which occurs adjacent to catalyst, can be thought of a hydrogen atom shift from the beta methylene to the alpha methylene converting it to the methyl branch. Catalyst (Fe, Co, Ru, etc.) moves from alpha to beta and with insertion of additional methylene(s) between catalyst and the methine group (former beta), isolation of the methyl branch is complete. The second key point is that alpha olefins can be a major product of F-T chemistry.

The present invention makes use of such observations to provide an overall method for preparing mid- or near-mid chain branched alpha-olefins which can be converted to the corresponding detergent surfactants, either directly or through the formation of intermediate compounds (e.g., branched-chain alcohols) which are subsequently converted into surfactants. Importantly, the surfactants thus made contain little or no contaminants such as the geminal or vicinal branches or multiple chain branches (i.e., more than about 3 branches). On a weight basis, such contaminants can detract from overall detergency performance and/or biodegradability of the final surfactant products herein.

The overall process herein is as follows:

The Fischer-Tropsch process is described in Kirk-Othmer Encyclopedia of Chemical Technology, 4th Edition, Volume 12, pp. 157-164 (1994), Jacqueline I. Kroschwitz, Executive Editor, Wiley-Interscience, N. Y. The Oxo process to make alcohols is described in detail in Kirk-Othmer Encyclopedia of Chemical Technology, 4th Edition, Volume 1, pp. 903-8 (1991).

1) Synthesis gas, a mixture of carbon monoxide/hydrogen is typically generated from coal or natural gas, however petroleum or other hydrocarbon sources could in principle be utilized. Air or oxygen is used to partially burn gas, petroleum, etc., to a mixture of carbon monoxide and hydrogen. Similarly, coal or coke can undergo the coke-water-gas reaction to form carbon monoxide and H₂. The water gas shift reaction can be used to change the carbon monoxide/hydrogen ratio as

WO 97/38956

PCT/US97/06339

5

required. Various standard cleanup steps are included to remove carbon dioxide, hydrogen sulfide, ammonia etc.

Gas + air or O₂ → CO/H₂ mixture

C + H₂O → CO + H₂ coke-water-gas reaction

5 CO + H₂O → H₂ + CO₂ (water gas shift)

2) Fischer-Tropsch (F-T) chemistry is used to convert synthesis gas into a mostly hydrocarbon mixture. Conditions can be set to produce a mostly linear olefin mixture with a limited number of methyl branches as well as some cyclic hydrocarbons. Small amounts of other classes of compounds such as alcohols are also
10 formed. Their levels can be somewhat controlled by F-T conditions; in any event they can be removed.

CO/H₂ → Syn Fuel Mixture + Branched Alpha-Olefins

3) Distillation and other standard techniques are used to isolate the desired MW hydrocarbon fraction containing alpha-olefins. Molecular sieving can be used to
15 separate most of the linear alpha-olefins and cyclics from the desired, limited methyl-branched, linear alpha-olefins. Standard methods utilizing zeolites can accomplish the former. Processing with zeolite sieves can be arranged to remove iso and anteiso (omega-1) and (omega-2) methyl alpha olefins, if so desired. Aliphatic hydrocarbons containing 2 geminal Me groups or highly branched aliphatic hydrocarbons (including
20 cyclics) can be separated from aliphatic hydrocarbons containing Me groups on different C atoms and less branched aliphatic hydrocarbons by selective adsorption of the latter on a molecular sieve (pore diam. 4.4-5.0Å) and/or from pyrolyzed poly(vinylidene chloride) (Saran) to yield gasoline with improved octane numbers; see Neth Appl. 7111508 10/25/71, Chem. Abstracts 76:88253.

25 Syn Fuel Mixture → Branched alpha-Olefins

4) Oxo chemistry (CO/H₂) is used to convert the branched alpha-olefin to the corresponding branched primary alcohol. Any Oxo catalyst which leads directly to alcohols or indirectly through an additional step of hydrogenation of intermediate aldehyde can be used. However it is preferable to use catalysts which do not
30 isomerize the double bond of the alpha-olefin prior to carbonylation as is the case using cobalt-carbon monoxide-organophosphine catalysts in the one step process. Conventional cobalt Oxo catalysts such as cobalt-carbon monoxide used in the two step high pressure process do not isomerize the C=C double bond. The fact these can give approximately equal carbonylation on 1- and 2-carbon positions of the alpha
35 olefin is entirely acceptable. In other words the product mixture would be RCH₂CH₂CH₂OH + RCH(CH₃)CH₂OH where R is linear fatty chain with limited methyl branching at the mid- or near-mid chain region.

WO 97/38956

PCT/US97/06339

6

Branched Alpha-olefins → Branched Primary Alcohols

- 5) In one aspect of the last step, any standard sulfation technique may be used to convert the above branched alcohol to a branched alcohol sulfate. Examples are sulfur trioxide in a falling film reactor or sulfur trioxide or chlorosulfonic acid in a batch reactor. In any case the acid mixture is promptly neutralized with caustic soda, or the like.

Branched Primary Alcohol → Branched Alkyl Sulfate.

- Other fatty alcohol-derived surfactants can also be made, e.g., alkyl ethoxyl sulfates (AES), alkyl polyglucosides (APG), etc. Note that surfactants other than alcohol sulfates or AES may be made by oxidizing said alcohol or its aldehyde intermediate into a carboxylate (i.e., a branched-chain soap). This soap can be an excellent surfactant and/or detergent builder in and of itself. This carboxylate can also be used as a feedstock and converted to branched acyl-auroates, -isethionates, -sarcosinates, -N-methylglucamides or other similar acyl-derived surfactants, using art-disclosed techniques.

INDUSTRIAL APPLICABILITY

- Branched-chain surfactants of the type resulting from the present process can be used in all manner of cleaning compositions. Such compositions include, but are not limited to: granular, bar-form and liquid laundry detergents; liquid hand dishwashing compositions; liquid, gel and bar-form personal cleansing products; shampoos; dentifrices; hard surface cleaners, and the like. Such compositions can contain a variety of conventional detergent ingredients. The following listing of such ingredients is for the convenience of the formulator, and not by way of limitation of the types of ingredients which can be used with the branched-chain surfactants herein.

- The branched-chain surfactants herein can be used in combination with detergency builders. Such builders include, for example, 1-10 micrometer zeolite A, polycarboxylate builders such as citrate, layered silicate builders such as "SKS-6" (Hoechst) and phosphate materials, especially sodium tripolyphosphate ("STPP"). Most laundry detergents typically comprise at least about 1% builder, more typically from about 5% to about 80% builder or mixtures of builders.

- Enzymes, such as proteases, amylases, lipases, cellulases, peroxidases, and mixtures thereof, can be employed in detergent compositions containing the branched-chain surfactants. Typical detergent compositions comprise from about 0.001% to about 5% of commercial enzymes.

Detergent compositions can also contain polymeric soil release agents (SRA's). Such materials include, for example, anionic, cationic and non-charged

WO 97/38956

PCT/US97/06339

7

monomer units, especially polyester materials. Preferred materials of this type include oligomeric terephthalate esters, sulfonated substantially linear ester oligomers comprising a backbone of terephthaloyl and oxyalkyleneoxy repeat units and phthalolyl-derived sulfonated terminal moieties. A variety of SRA's are described, for example, in U.S. 4,968,451; 4,711,730; 4,721,580; 4,702,857; 4,877,896; 5,415,807; and in other literature references. Such soil release materials typically comprise from about 0.01% to about 10% of finished detergent compositions.

Detergent compositions may also optionally contain bleaching compositions comprising a bleaching agent and one or more bleach activators. If present, bleaching agents such as percarbonate or perborate (especially perborate monohydrate "PB1") typically are used at levels from about 1% to about 30% of finished detergent compositions. Bleach activators such as nonanoyloxy-benzene sulfonate ("NOBS") and tetraacetyl ethylenediamine ("TAED"), and mixtures thereof, can be used to enhance the bleaching activity of materials such as perborate and percarbonate. If present, the amount of bleach activator will typically be from about 0.1% to about 60% of a bleaching composition comprising a bleaching agent-plus-bleach activator. Other bleaching agents such as the so-called "photoactivated" bleaches (see U.S. 4,033,718) can also be used. Sulfonated zinc phthalocyanine is an especially preferred photoactivated bleaching agent.

Detergent compositions can also contain clay soil removal/antiredeposition agents such as ethoxylated tetraethylene pentamine; see U.S. 4,597,898. Such materials typically comprise from about 0.01% to about 10% of fully-formulated laundry detergents.

Detergent compositions can also contain from about 0.1% to about 7% of polymeric dispersing agents, which are especially useful in the presence of zeolite and/or layered silicate builders. Such materials are known in the art (see U.S. 3,308,067). Such materials include acrylate/malic-based copolymers, such as described in EP 193,360, as well as polyethylene glycol ("PEG").

Detergent compositions herein can also include various brighteners, dye transfer inhibiting agents (especially polymers of N-vinylpyrrolidone and N-vinylimidazole), suds suppressors (especially silicones), chelating agents such as nitrilotriacetate, ethylenediamine disuccinate, and the like. Such materials will typically comprise from about 0.5% to about 10%, by weight, of fully-formulated cleaning compositions.

Moreover, it is to be understood that the branched-chain surfactants prepared in the manner of the present invention may be used singly in cleaning compositions or in combination with other deterative surfactants. Typically, fully-formulated cleaning

WO 97/38956

8

PCT/US97/06339

compositions will contain a mixture of surfactant types in order to obtain broad-scale cleaning performance over a variety of soils and stains and under a variety of usage conditions. One advantage of the branched-chain surfactants herein is their ability to be readily formulated in combination with other known surfactant types. Nonlimiting examples of additional surfactants which may be used herein typically at levels from about 1% to about 55%, by weight, include the unsaturated sulfates such as oleyl sulfate, the C₁₀-C₁₈ alkyl alkoxy sulfates ("AE_xS"; especially EO 1-7 ethoxy sulfates), C₁₀-C₁₈ alkyl alkoxy carboxylates (especially the EO 1-5 ethoxycarboxylates), the C₁₀-18 glycerol ethers, the C₁₀-C₁₈ alkyl polyglycosides and their corresponding sulfated polyglycosides, and C₁₂-C₁₈ alpha-sulfonated fatty acid esters. Nonionic surfactants such as the ethoxylated C₁₀-C₁₈ alcohols and alkyl phenols, (e.g., C₁₀-C₁₈ EO (1-10) can also be used. If desired, other conventional surfactants such as the C₁₂-C₁₈ betaines and sulfobetaines ("sultaines"), C₁₀-C₁₈ amine oxides, and the like, can also be included in the overall compositions. The C₁₀-C₁₈ N-alkyl polyhydroxy fatty acid amides can also be used. Typical examples include the C₁₂-C₁₈ N-methylglucamides. See WO 9,206,154. Other sugar-derived surfactants include the N-alkoxy polyhydroxy fatty acid amides, such as C₁₀-C₁₈ N-(3-methoxypropyl) glucamide. The N-propyl through N-hexyl C₁₂-C₁₈ glucamides can be used for low sudsing. C₁₀-C₂₀ conventional soaps may also be used. If high sudsing is desired, the branched-chain C₁₀-C₁₆ soaps may be used. C₁₀-C₁₄ alkyl benzene sulfonates (LAS), which are often used in laundry detergent compositions, can also be used with the branched surfactants herein.

The following Examples illustrate the use of branched-chain surfactants prepared according to the present invention in various cleaning compositions, but is not intended to be limiting thereof.

EXAMPLE I

Granular laundry detergents are prepared as follows.

		A	B	C
	Blown Powder			
30	Zeolite A	30.0	22.0	6.0
	Sodium sulfate	19.0	5.0	7.0
	Polyacrylate			
	LAS	13.0	11.0	21.0
	Branched AS*	9.0	8.0	8.0
35	Silicate, Na	-	1.0	5.0
	Soap	-	-	2.0
	Carbonate, Na	8.0	16.0	20.0

WO 97/38956

PCT/US97/06339

9

Spray On				
	C ₁₄₋₁₅ EO7	1.0	1.0	1.0
Dry additives				
5	Protease	1.0	1.0	1.0
	Lipase	0.4	0.4	0.4
	Amylase	0.1	0.1	0.1
	Cellulase	0.1	0.1	0.1
	NOBS	-	6.1	4.5
	PB1	1.0	5.0	6.0
10	Sodium sulfate	-	6.0	-

Moisture & Miscellaneous

Balance

*C₁₂-C₁₄ methyl branched alkyl sulfate, prepared as disclosed above.

WO 97/38956

10

PCT/US97/06339

A bleach-containing nonaqueous liquid laundry detergent is prepared as follows.

EXAMPLE II

	<u>Component</u>	<u>Wt. %</u>	<u>Range (% wt.)</u>
5	<u>Liquid Phase</u>		
	Branched AS*	25.3	18-35
	C ₁₂₋₁₄ EO5 alcohol ethoxylate	13.6	10-20
	Hexylene glycol	27.3	20-30
	Perfume	0.4	0-1.0
10	<u>Solids</u>		
	Protease enzyme	0.4	0-1.0
	Na ₃ Citrate, anhydrous	4.3	3-6
	Sodium perborate (PB-1)	3.4	2-7
	Sodium nonanoyloxybenzene sulfonate (NOBS)	8.0	2-12
15	Sodium carbonate	13.9	5-20
	Diethyl triamine pentaacetic acid (DTPA)	0.9	0-1.5
	Brightener	0.4	0-0.6
	Suds Suppressor	0.1	0-0.3
	Minors	—————Balance—————	
20	*C ₁₂ -C ₁₆ methyl branched alkyl sulfate, Na salt, prepared as disclosed above.		

A hand dishwashing liquid is as follows.

EXAMPLE III

	<u>Ingredient</u>	<u>% (wt.)</u>	<u>Range (% wt.)</u>
25	Branched AS*	13.0	5-15
	Ammonium C ₁₂₋₁₃ alkyl ethoxy sulfate	15.0	10-35
	Cocnut amine oxide	2.6	2-5
	Betaine**/Tetronic 704®	0.87-0.10	0-2 (mix)
30	Alcohol Ethoxylate C ₈ E ₁₁	5.0	2-10
	Ammonium xylene sulfonate	4.0	1-6
	Ethanol	4.0	0-7
	Ammonium citrate	0.06	0-1.0
	Magnesium chloride	3.3	0-4.0
35	Calcium chloride	2.5	0-4.0
	Ammonium sulfate	0.08	0-4.0
	Hydrogen peroxide	200 ppm	0-300 ppm

WO 97/38956

PCT/US97/06339

11

Perfume	0.18	0-0.5
Maxatase® protease	0.50	0-1.0
Water and minors	Balance	

5 *C₁₂C₁₄ methyl branched alkyl sulfate, triethanolammonium salt, prepared as disclosed above.

**Cocoalkyl betaine.

WO 97/38956

PCT/US97/06339

12

WHAT IS CLAIMED IS:

1. A process for preparing mid- to near-mid chain branched alpha olefins, comprising the steps of:

- (a) preparing a mixture of CO and H₂;
- (b) reacting the mixture of CO and H₂ in the presence of a catalyst under Fischer-Tropsch conditions to prepare a hydrocarbon mixture comprising said mid- to near-mid chain branched alpha-olefins; and
- (c) separating said branched alpha-olefins from said hydrocarbon mixture.

2. A process according to Claim 1 wherein the catalyst of step (b) is a member selected from the group consisting of Fe, Co and Ru Fischer-Tropsch catalysts.

3. A process according to Claim 1 wherein said alpha-olefins are separated using molecular sieves and/or pyrolyzed poly(vinylidene chloride).

4. A process for preparing mid- to near-mid chain branched alcohols, comprising reacting the branched alpha-olefins made in accordance with Claim 1 with CO/H₂ under Oxo conditions.

5. A process according to Claim 4 which is conducted without isomerization of the olefin double bond using a cobalt-carbon monoxide catalyst.

6. A process according to Claim 5 which yields alcohols of the type RCH₂CH₂CH₂OH and RCH(CH₃)CH₂OH, wherein R is C₈ to C₁₆ alkyl with methyl branching.

7. A process for preparing branched alkyl sulfate surfactants, comprising sulfating the alcohols prepared according to Claim 4.

8. A process for preparing branched alkyl ethoxy sulfates, comprising first ethoxylating, then sulfating, the alcohols prepared according to Claim 4.

9. A process for preparing branched alkyl carboxylate surfactants by oxidizing the alcohols prepared according to Claim 4, or their aldehyde intermediates.

WO 97/38956

PCT/US97/06339

13

10. A process for preparing branched acyl taurate, branched acyl isethionates, branched acyl sarcosinate or branched acyl N-methylglucamide surfactants, or the like, using the branched carboxylates prepared according to Claim 9 as a feedstock.

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.
PCT/US 97/06339

A. CLASSIFICATION OF SUBJECT MATTER		
IPC 6	C07C1/04 C07C305/06	C07C11/02 C07C305/10
	C07C29/16 C07C51/16	C07C31/125 C11D1/14
C07C303/24		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
IPC 6 C07C		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	H.F. MARK, ET AL.: "Kirk-Othmer Encyclopaedia of Chemical Technology, 3rd. edition, vol. 11" 1989, JOHN WILEY & SONS, NEW YORK, US XP002036088 pages 447 - 489 see page 473 - page 476 ---	1
A	H.F. MARK, ET AL.: "Kirk-Othmer Encyclopaedia of Chemical Technology, 3rd. edition, vol. 16" 1989, JOHN WILEY & SONS, NEW YORK, US XP002014913 see page 637 - page 653 ---	4
-/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "A" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
25 July 1997		05. 08. 97
Name and mailing address of the ISA European Patent Office, P.B. 3818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+ 31-70) 340-3016		Authorized officer English, R

Form PCT/ISA/216 (second sheet) (July 1992)

page 1 of 2

INTERNATIONAL SEARCH REPORT

Intern. Application No
PCT/US 97/06339

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3 480 556 A (W.J. DE WIT, ET AL.) 25 November 1969 cited in the application see the whole document -----	7,8

1

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

page 2 of 2

INTERNATIONAL SEARCH REPORT
Information on patent family membersIntern al Application No
PCT/US 97/06339

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 3480556 A	25-11-69	NONE	

Form PCT/ISA/210 (patent family annex) (July 1992)